#### HIV是什麽

#### What is HIV?

HIV是一種感染人體免疫系統的病毒,它會破壞或 菌會趁免疫系統虛弱的時候感染人體 造成相關感染稱為「伺機性感染」,原因為其他病 病感染時,就稱為「免疫缺乏」。嚴重的免疫缺乏 系統逐漸惡化,當免疫系統惡化到無法抵抗其他疾 損傷免疫系統的功能。感染HIV後會造成人體免疫

associated with severe immunodeficiency are deficiency". The immune system is considered known as "opportunistic infections", because deficient when it can no longer fulfill its role of of the immune system, leading to "immune with the virus results in progressive deterioration destroying or impairing their function. Infection The HIV infects cells of the immune system system they take advantage of a weakened immune fighting infection and disease. Infections

### 正確使用男性保險套

Five steps for Correct

25大步飄

Sharing needles, syringes or solution

母子垂直傳染 性行為

use of male condoms



Tear 期






#### HIV如何傳染

## How is HIV fransmitted?

childbirth and breastfeeding contaminated needles, syringes or other sharp contaminated blood; and the sharing of sex with an infected person; transfusion of sexual intercourse [ vaginal or anal ], and oral HIV can be transmitted through unprotected 共用針頭、稀釋液或其他尖銳器具皆是感染途徑 HIV可以透過無保險套的性行為(包括陰道性交或 instruments. It may also be transmitted between 口交)以及肛交來感染;輸入感染HIV的血液, 它也可能透過女性懷孕、分娩及哺乳間傳染給嬰兒 mother and her infant during pregnancy. 及







## 如何避免自己暴露在HIV

感染風險中

#### transmission through sex? How can I minimize my risk of HIV

- (1)在每次性行為時都可以正確的使用保險套 (包括無套肛交、性交及口交)
- (2)保險套要選擇水性潤滑液來搭配
- (3)限制你的性伴侶人數。
- (4)若你認為自己有可能感染HIV,請儘快在暴 露後72小時內至醫院找醫師討論是否該進行 預防性投藥治療
- (5)若你的性伴侶HIV呈現陽性,鼓勵他立刻開 始並持續治療。
- (1) Use condoms correctly each time you have sex (include anal sex, vaginal sex and oral sex).
- (2) Choose water-based lubricants for use with condoms
- (3) Limit your number of sex partners
- (4) Talk to your doctor right away (within 3 days) have a possible exposure to HIV about post-exposure-prophylaxis (PEP) if you
- (5) If your partner is HIV-positive, encourage your partner to get and stay on treatment



#### TESTING



# What is HIV testing?

都可以因篩檢使自己和親人有更好的保護。 篩檢是一個可以讓你了解HIV感染狀況的有效工具, 是免費、快速且保密的。篩檢後不管結果如何,你 能夠讓你知道自己是否有感染HIV就只有篩檢。篩檢 你可能已經感染HIV了但你卻認為現在很健康。因此

and your loved ones. use that knowledge to take better care of yourself status. You could have HIV and still feel healthy. HIV testing is an effective tool to know your HIV Whether you test positive or negative, you can tested. HIV testing is free, fast, and confidential The only way to know if you have HIV is to get

### 何時該做HIV篩檢

# When should I get tested for HIV?

持續6-12週。因此若你覺得自己可能有感染HIV,建議 抗體尚未產生,此期間稱為空窗期,愛滋病空窗期約 在感染HIV初期,篩檢的結果可能是陰性,是因為HIV 你在暴露後6-12週內去做HIV篩檢最準確。

approximately 6 to 12 weeks. If you think you may period of time is called the window period. It lasts because antibodies have not been produced. This testing 6 to 12 weeks after exposure have been exposed to HIV, we recommend HIV At the early infection of HIV, test may be negative



Time/ 13:00~20:00 (每週一/國定假日-休館 Closed on Mondays and national holidays

e-Mail/ rainbowst6@gmail.com Tel/ (06)263-1841 · 263-7641

Add/70261台南市南區南和路 6號3樓 3F., No.6, Nanhe Rd., South Dist., Tainan City 70261, Taiwar (台南市南都里活動中心對面)

Located opposite to the Nandu Village Community Center



FB粉絲專頁

#### 臺南市政府衛生局

請於上班時間洽詢

東興辦公室) 06-633-3080・(林森辦公室) 06-336-6366

麥滋病免費諮詢專線:1922

愛滋病免費篩檢:臺南市政府衛生局及各區衛生所 Free HIV Test : Public Health Bureau, Tainan City Gov

各大醫院愛滋病匿名篩檢電話:

成大醫院門診大樓1樓 (06)235-3535轉4100 First floor at the Outpatient Building, National Cheng Kung University Hospital

奇美醫院永康院區2樓抽血站 (06)2812811轉53689 Drawing Station on the second floor, Chi Mei Hospital, YongKang

衛生福利部臺南醫院門診大樓2樓感控室 (06)220-0055轉3125

http://www.cdc.gov.tw/rwd ion Control Office on the Second floor at the Outpatient Building,

衛生福利部疾病管制署民眾版



www.cdc.gov.tw



1922防疫達人 www.facebook.com/TWCDC

迅

3

这

查 疤


H

匨

过



LOVE YOURSELF, LOVE OTHERS



衛生幅利部疾病管制署

图 關心您 Sponsored by

廣告 AD